

Short Communication

Ethnic Differences in the Prevalence, Clinical Outcome and *cag* Pathogenicity Island (*cagPAI*) Virulence Gene Profiles of *Helicobacter pylori* Strains from Malaysia

Hamat, R. A.^{1*}, Nor Amalina, E.², Malina, O.¹, Zamberi, S.¹, Alfizah, H.³, Rizal, A. M.⁴, Aminuddin, A.⁵ and Ramelah, M.⁶

¹Department of Medical Microbiology and Parasitology, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

²School of Medicine, University Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu, Sabah, Malaysia

³Department of Medical Microbiology and Immunology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, 56000 Cheras, Kuala Lumpur, Malaysia

⁴Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, 56000 Cheras, Kuala Lumpur, Malaysia

⁵Faculty of Medicine, Universiti Teknologi MARA Selayang Campus, Jalan Prima Selayang 7, 68100 Batu Caves, Selangor, Malaysia

⁶Collaborative Innovation Centre, Universiti Kebangsaan Malaysia, Bangi, 43600 Bangi, Selangor, Malaysia

ABSTRACT

Different *Helicobacter pylori* genes may be well conserved within different ethnic groups and could give rise to different clinical outcomes. In this study, we demonstrated a low prevalence of *H. pylori* infection (19.2%) which is in concordance with the current trend demonstrated locally and abroad. The Indians had the highest prevalence of *H. pylori* infection among other ethnic groups (Malays= 8.6 %, Chinese= 24.3 %, Indians= 33.9%). *cagM* and *cagT* were the most predominant genes found (63.4% for each), followed by *cagA* (62.2 %), *cagE* (48.2%), *cag6-7* (46.3%), *cagI0* (42.1%), *cagI3* (4.9%) and *IS605* (3.7%). No significant association was found between *H. pylori* infection and *H. pylori* genes

with ethnic groups or clinical outcomes. Indians who had a combination of *cagA*/*E/M* genes of *H. pylori* were likely to be associated with 21-time of having non-ulcer dyspepsia (NUD) than peptic ulcer disease (PUD). Therefore, these genes may serve as useful markers in predicting the clinical presentation of a *H. pylori* infection among

ARTICLE INFO

Article history:

Received: 2 April 2012

Accepted: 6 May 2013

E-mail addresses:

rukman@upm.edu.my (Hamat, R. A.),
e.amalina@hotmail.com (Nor Amalina, E.),
malinaosman@upm.edu.my (Malina, O.),
zamberi@upm.edu.my (Zamberi, S.)
alfizah@ppukm.ukm.my (Alfizah, H.),
mrizal@ppukm.ukm.my (Rizal, A. M.),
ramelahm@yahoo.com (Ramelah, M.)

* Corresponding author

Indians in our studied population. Hence, this preliminary data might explain why Indians have a low prevalence of gastric cancer and peptic ulcer disease despite having persistently high prevalence of *H. pylori* infection for many decades (“Indian enigma”) in Malaysian patients.

Keywords: *Helicobacter pylori*, *cag* pathogenicity island, virulence genes, peptic ulcer diseases, non-peptic ulcer dyspepsia, ethnicity, Malaysia

INTRODUCTION

Helicobacter pylori is present in more than 50% of the world’s population (Peterson, 1992). This unique bacterial species is extremely diverse in its genomic structures. This panmicticism is believed to be important in colonization and infection in the human gastro-duodenal system. A variety of diseases including gastritis, gastric ulcers (GU), duodenal ulcers (DU) and gastric cancer have been attributed to *H. pylori*. The prevalence rates of *H. pylori* vary between populations or sub-populations (ethnic groups) within the same geographic locations (Suerbaum & Michetti, 2002). Data from endoscopic surveys and sero-prevalence studies amongst the three Malaysian ethnic groups revealed that Indians have the highest prevalence of *H. pylori* infection (68.9% -75.0%) compared to Chinese (45.0% - 60.6%), while Malays consistently have the lowest prevalence rate (8.0% - 43.3%). Ironically, in terms of the severity of disease caused by *H. pylori*, gastric cancer occurs more frequently in Chinese (68.0%) rather than Indians

(16.5%) and Malays (15.5%)(Haron *et al.*, 1994; Kang *et al.*, 1990). In addition, it has been documented that peptic ulcer disease (PUD) is rather low among Indians compared to Chinese (Owen *et al.*, 2001; Goh, 1997). This paradox which is known as the “Indian enigma” (high prevalence of *H. pylori* but low percentage of gastric cancer and PUD) may be explained by differential virulence of the *H. pylori* strains or different susceptibility genotypes in the Indian ethnic group. *cagPAI*, a 40-kb region of first chromosomal DNA is the major genetic determinant of *H. pylori* virulence that consists of 27 genes and it is divided into two segments, *cag* I and *cag* II, by an insertion sequence known as *IS605* (Censini *et al.*, 1996). There is now increasing evidence for the existence of several novel bacterial pathogenicity markers on the *cag* (cytotoxin-associated gene) pathogenicity island (*cagPAI*) that may play important roles in the disease invoking potential of *H. pylori* (Mattar *et al.*, 2007; Van Doorn *et al.*, 1998). Meanwhile, the overall genetic variability of *H. pylori* may also be held responsible for the great diversity of clinical outcomes (Proença-Módena *et al.*, 2007; Maeda *et al.*, 1999). However, this has not been well explored locally in Malaysia and detailed data is lacking. Therefore, we conducted a study to investigate the role of several selected genes in the *cagPAI* (*cagA*, *cagE*, *cagM*, *cagT* and *cag6-7*, *cag10*, *cag13* represented the *cag* I and *cag* II segments, respectively and *IS605*) versus ethnicity and clinical outcomes. To our knowledge, this is the largest set of *cagPAI* genes that has

been studied so far in *H. pylori* from clinical isolates in Malaysia.

MATERIALS AND METHODS

A total of 855 gastric biopsies were obtained from dyspeptic patients who had undergone endoscopy from May 2004 to May 2007 in Universiti Kebangsaan Malaysia Medical Center. These patients were enrolled by a purposive sampling. Selection of patients and study protocols were followed according to a previous local study (Ramelah *et al.*, 2005). The study protocol was approved by the Medical Ethics Committee of Universiti Kebangsaan Malaysia (FF-075-2003). *H. pylori* culture was performed as previously described with modifications (Proença-Módena *et al.*, 2007). Briefly, specimens for culture were immediately transported to the laboratory in Stuart transport medium (Oxoid, UK). Biopsies were sub-cultured onto Columbia agar base medium (Oxoid Ltd., Basingstoke, Hampshire, England) containing Dent's supplement and 7% lysed horse blood. The cultures were incubated under micro-aerophilic conditions (5–6% O₂, 8–10% CO₂, 80–85% N₂, and a relative humidity of at least 95%) at 37°C for a minimum period of 5 days. The bacteria were identified as *H. pylori* based on the typical Gram stain morphology and biochemical tests such as urease, oxidase and catalase reactions. A pool of bacterial colonies obtained from each single plate was used for DNA extraction. Two *H. pylori* reference strains: American type culture collection (ATCC) 700824 (strain J99) and ATCC 43256 were

used in this study. Chromosomal DNA of each strain was prepared using the High Pure PCR Template Preparation kit (Roche, Mannheim, Germany) in accordance with the manufacturer's instructions.

PCR analysis for *cagPAI* was performed to amplify *cagA*, *cagE*, *cagM*, *cagT*, *cag6-7*, *cag10*, *cag13* and *IS605* genes. The primers and PCR conditions used for each gene have been reported previously by Maeda *et al.* (1999). Briefly, the PCR was performed in a total reaction volume of 25 µl containing 2.5 µl 10 X PCR buffer, 1 µl of each primer (each at 10 pmol/µl), 2 mM MgCl₂, 200 µM of dNTP, 5 U of super Taq (Super Taq DNA Polymerase Mbiotech) and 1 µL (10 ng) of genomic DNA from the culture lysates. DNA extracts of *H. pylori* ATCC 700824 and *H. pylori* ATCC 43256 were used as positive controls while negative controls (without DNA) were added to each PCR run. PCR products were visualized by electrophoresis in 1.2% agarose gel, stained with ethidium bromide, and examined under UV illumination. Standard 1-kb DNA ladders (Fermentas Life Science, Hanover, MD) were used as molecular size markers.

Chi-squared and Fisher's exact tests were used to determine the difference in the distribution of *H. pylori* genes and the correlation between genes, clinical disease and ethnicity. *p* values of less than 0.05 were considered statistically significant.

RESULTS AND DISCUSSION

In this study, we reported a lower overall prevalence of *H. pylori* amongst our studied population (19.2%; 164 out of 855) compared

to 49% and 22% by Goh *et al.* (1997) and Mahadeva *et al.* (2005), respectively. This is not surprising as high standard of living conditions and improved personal hygiene could contribute to major decline of *H. pylori* infection in Asian populations (Tan & Goh, 2008). Interestingly, our finding corroborates data from a study conducted in a different locality, i.e. the north eastern part of Peninsular Malaysia which was 19.3% (Raj *et al.*, 2001). With regard to ethnic differences, a similar pattern was observed; Indians had the highest prevalence rate of *H. pylori* infection (33.9%; 42/124) compared to Chinese (24.3%; 98/404) and Malays (8.6%; 28/327) (data not shown). This is very much consistent with previous local findings as well (Mahadeva *et al.*, 2005; Goh *et al.*, 1997; Kang *et al.*, 1990) and could be explained by the role of genetic heterogeneity in susceptible hosts or other co-factors that could possibly account for the inter-ethnic variations (Alm & Trust, 1999).

For instance, *cagA* gene in *cagPAI* region of *H. pylori* strains was responsible for more severe pathological changes and clinical outcomes in Western countries (Crabtree *et al.*, 1993; Xiang *et al.*, 1993). Surprisingly, similar correlations were not found in some Asian countries (Zheng *et al.*, 2000; Maeda *et al.*, 1998). *cagA* gene was reported to be insufficient in inducing some disease processes in *H. pylori* infection in China (Hu *et al.*, 1995). Tan *et al.* (2005) reported 84% of their patients who had *cagA* gene were not associated with clinical outcomes. In addition, no

significant correlation was observed in the prevalence of this gene among the Malays (76.6%), Chinese (86.4%) and Indians (86.8%). In our previous report, we could not reveal the association between either the *cagA* or 3' end region of *cagA* gene with gastroduodenal diseases. However, Chinese (96%) in whom have been known to have a higher risk of peptic ulcer disease and gastric cancer had the highest prevalence of *cagA* subtype A strains compared to Malays (72%) and Indians (69%). The difference was statistically significant ($p < 0.0005$) (Ramelah *et al.*, 2005). In a recent work by our group, Eastern CagA strains were significantly predominance among the Chinese (82.9%) compared to Malays (11.4%) and Indians (1.4%). However, no association could be made between Eastern CagA strains and the clinical outcome (Mohamed *et al.*, 2009). It has been known that Eastern CagA strains confer higher tyrosine CagA phosphorylation activity than Western strains, which is related to their pathogenic potentials (Higashi *et al.*, 2002a). As previous works were only focussing on *cagA* and its variants, and could not demonstrate the association with clinical outcome, we hypothesized that other potential virulence genes on *cagPAI* could also be involved in *H. pylori* infection among our multi-ethnic population. Maeda *et al.* (1999) evaluated 15 genes on the *cagPAI* and concluded that *cagA* gene could not be used as a virulence marker in Japanese *H. pylori* strains. Recently, a systematic mutagenesis study of individual genes on *cagPAI* by Fischer *et al.* (2001)

confirmed that 17 and 14 out of 27 genes were involved in *H. pylori* pathogenesis by encoding a type IV secretion system for the translocation of *cag* toxicity protein A (CagA) and for the full induction of interleukin 8 (IL-8). Among the eight genes, *H. pylori cagM* and *cagT* genes were both predominantly found in our study population (63.4% for each) (TABLE 1). To

the best of our knowledge, this is the first report on the prevalence of *cagM* and *cagT* genes in Malaysia; nonetheless, we could not reveal any significant association in the prevalence of *cagM* and *cagT* genes as well as other *cag*PAI genes with clinical outcome (TABLE 2). Goh *et al.* (2007) suggested that other virulent genes could account for the paradoxical findings of *H. pylori* infection

TABLE 1

Distribution of the presence of *cag*PAI genes of *Helicobacter pylori* (*cagA*, *cagE*, *cagM*, *cagT*, *cag6-7*, *cag10*, *cag13* and *IS605*) detected in 164 patients in relation to their ethnicity

Gene	Ethnic group			Total
	Malay n = 28 (%)	Chinese n = 94 (%)	Indian n = 42 (%)	
<i>CagA</i> (n=102)				
Positive	16 (57.1)	62 (66.0)	24 (57.1)	102 (62.2)
Negative	12 (42.9)	32 (34.0)	18 (42.9)	62 (37.8)
<i>CagE</i> (n=102)				
Positive	13 (46.4)	44 (46.8)	22 (52.4)	79 (48.2)
Negative	15 (53.6)	50 (53.2)	20 (47.6)	85 (51.8)
<i>CagM</i> (n=102)				
Positive	16 (57.1)	59 (62.8)	29 (69.0)	104 (63.4)
Negative	12 (42.9)	35 (37.2)	13 (31.0)	60 (36.6)
<i>CagT</i> (n=102)				
Positive	16 (57.1)	62 (66.0)	26 (61.9)	104 (63.4)
Negative	12 (42.9)	32 (34.0)	16 (38.1)	60 (36.6)
<i>Cag6-7</i> (n=102)				
Positive	8 (28.6)	59 (62.8) ^a	9 (21.4)	76 (46.3)
Negative	20 (71.4)	35 (37.2)	33 (78.6)	88 (53.7)
<i>Cag10</i> (n=102)				
Positive	12 (42.9)	38 (40.4)	19 (45.2)	69 (42.1)
Negative	16 (57.1)	56 (59.6)	23 (54.8)	95 (57.9)
<i>Cag13</i> (n=102)				
Positive	2 (7.1)	3 (3.2)	3 (7.1)	8 (4.9)
Negative	26 (92.9)	91(96.8)	39 (92.9)	156 (95.1)
<i>IS605</i> (n=102)				
Positive	0 (0.0)	3 (3.2)	3 (7.1)	6 (3.7)
Negative	28 (100.0)	91 (96.8)	39 (92.9)	158 (96.3)

^aChinese vs. Malay (Pearson Chi-square, 2 x 2 table; $\chi^2 = 10.189$, $p = 0.001$)

Chinese vs. Indian (Pearson Chi-square, 2 x 2 table; $\chi^2 = 19.842$, $p = 0.000$)

TABLE 2

Relationship between *Helicobacter pylori* genes (*cagA*, *cagE*, *cagM*, *cagT*, *cag6-7*, *cag10*, *cag13* and *IS605*) and clinical outcomes in 164 patients with *H. pylori* infection

Gene	PUD n=35 (%)	NUD n=129 (%)	Adjusted OR (95 % CI)	p value ^a
<i>cagA</i> (n=102)	19 (54.3)	83 (62.2)	0.658 (0.309-1.402)	0.327
<i>cagE</i> (n=79)	14(40.0)	65 (48.2)	0.656 (0.307-1.403)	0.341
<i>cagM</i> (n=104)	22 (62.9)	82 (63.6)	0.970 (0.447-2.103)	1.000
<i>cagT</i> (n=104)	21 (60.0)	83 (64.3)	0.831 (0.386-1.789)	0.694
<i>cag6-7</i> (n=76)	17 (48.6)	59 (45.7)	1.121 (0.530-2.367)	0.849
<i>cag10</i> (n=69)	16 (45.7)	53 (42.1)	1.208 (0.569-2.561)	0.623
<i>cag13</i> (n=8)	2 (5.7)	6 (4.7)	1.242 (0.240-6.442)	0.679
<i>IS605</i> (n=6)	1 (2.9)	5 (3.9)	0.157 (0.082-6.455)	1.000

Note: PUD = peptic ulcer disease; NUD = non-ulcer dyspepsia; OR = odds ratio; CI = confidence interval

^aFischer's Exact test; p value < 0.05 is considered significant

and clinical outcome in our multi-ethnic groups. They found that only 23.3% of *H. pylori*-positive Indians had gastric cancer compared to 82.3% and 60.5% in Malays and Chinese, respectively. Mohamed *et al.* (2009) reported Western CagA strains were significantly found in Indian patients (43.5%), which could probably explained lower incidence of gastric cancer or PUD in this ethnic group. In our study, when each ethnic group was analyzed, the association was only statistically significant in the proportion of positive *cagA/E/M* genes amongst Indians who had NUD than PUD ($p < 0.047$) (TABLE 3). In addition, Indians with the combination of *cagA/E/M* genes were likely (twenty-one times) associated with NUD than PUD. These findings are rather surprising as the presence of these markers has been associated with high risk of developing PUD and bad clinical outcome (Mattar *et al.*, 2007; Censini *et al.*, 1996). *cagA* gene in particular was predominantly found in patients with DU in

both European and Polynesian in Auckland (83% and 86%, respectively) (Campbell *et al.*, 1997). Meanwhile, *cagE* gene was proposed to be a more reliable virulence marker than *cagA* gene (Ikenoue *et al.*, 2001; Audibert *et al.*, 2001; Maeda *et al.*, 1999) but Tan *et al.* (2005) reported that Chinese (39.0%) had significantly lower prevalence rate of *cagE* gene than Malay (70.0%) and Indian (81.6%) patients which appears *cagE* gene is not a good marker. *cagE* and *cagM* genes are responsible for activating the transcription factor NF- κ B, which mediates IL-8 secretion (Glocker *et al.*, 1998). However, our findings corroborate data with a study conducted in Taiwan which revealed no association between the presence of *cagA*, *cagE* and *cagM* genes with the type of disease and/or the histological findings in their patients (Sheu *et al.*, 2002). Furthermore, the involvement of these genes and others in *H. pylori* pathogenesis has also been disputed (Hsu P-I, *et al.*, 2002; Segal *et al.*, 1999). In

TABLE 3

Relationship between *Helicobacter pylori* genes (*cagA*, *cagE*, *cagM*, *cagT*, *cag6-7* and *cag10*) and clinical outcomes in 42 Indian patients with *H. pylori* infection

Gene (s)	PUD n=7 (%)	NUD n=35 (%)	Adjusted OR (95 % CI)	p value
<i>cagA</i> (n=24)	1 (4.2)	23 (95.8)	0.087 (0.009-0.808)	0.031 ^a
<i>cagE</i> (n=22)	1 (4.5)	21 (95.5)	0.111 (0.012-1.025)	0.041 ^b
<i>cagM</i> (n=29)	2 (6.9)	27 (93.1)	0.119 (0.019-0.731)	0.021 ^c
<i>cagT</i> (n=26)	3 (11.5)	23 (88.5)	0.391 (0.075-2.041)	0.397
<i>cag6-7</i> (n=9)	1 (11.1)	8 (88.9)	0.562 (0.059-5.386)	1.000
<i>cag10</i> (n=19)	1 (5.3)	18 (94.7)	0.157 (0.051-1.447)	0.105
<i>cagA/E/M</i> (n=21)	1 (4.8)	20 (95.2)	0.125 (0.014-1.151)	0.047 ^d

Note: PUD = peptic ulcer disease; NUD = non-ulcer dyspepsia; OR = odds ratio; CI = confidence interval
Statistical analysis was not done for *cag13* and *IS605* genes as number of samples was too small

^{a,b,c,d} by Fischer's exact tests; p values <0.05 are considered significant

addition, the prevalence of *cag6-7* gene in this study was significantly higher amongst the Chinese (62.8 %) compared to the Malays (28.6 %) (TABLE 1). The difference in the prevalence of *cag6-7* gene was highly statistically significant between Chinese and Indians (p=0.000). Interestingly, a study in Japan revealed that 93.1% (27/29) of cancer patients had *cag6-7* gene although the significance of this is still uncertain (Deguchi *et al.*, 2004), as this finding was not analyzed statistically.

Our study had several limitations. All *cagPAI* genes could not be chosen for the study due to budget constraints. The number of Indian patients was relatively small despite the significant findings observed in this ethnic group. However, the present study was conducted as a pilot study involving the largest number of *H. pylori* genes in different ethnic groups in Malaysia. Thus, large prospective or multi-centered studies are needed to investigate the pathogenic impact of *H. pylori* virulence genes amongst

different ethnic groups in relation to its clinical relevance.

CONCLUSION

The decline in the prevalence of *H. pylori* infection has not only been observed in several developed countries but also in Malaysia. Detection of *cagA/E/M* genes will partially resolve the "Indian enigma": it might be that the presence of different *H. pylori* genes/genotypes that might explain why Indians have a lower risk of developing severe disease outcomes despite having the highest prevalence rate of *H. pylori* infection. This could have clinical implication when initiating anti-*H. pylori* therapy in the multi-ethnic population in Malaysia.

ACKNOWLEDGMENTS

This study was supported by a grant from the Ministry of Science, Technology and Innovation of Malaysia (No. 06-02-04-0907-PR0073/05-2). Our special thanks

go to the staff of the Endoscopy and Histopathology Unit, Universiti Kebangsaan Malaysia Medical Center for their laboratory assistance. We also thank The Dean of Faculty of Medicine and Health Sciences, Universiti Putra Malaysia for her support as well.

REFERENCES

- Alm, R. A., & Trust, T. J. (1999). Analysis of the genetic diversity of *Helicobacter pylori*: the tale of two genomes. *The Journal of Molecular Medicine*, 77, 834–846.
- Audibert, C., Burucoa, C., Janvier, B., & Fauchere, J. L. (2001). Implication of the structure of the *Helicobacter pylori* *cag* pathogenicity island in induction of interleukin-8 secretion. *Infection and Immunity*, 69, 1625–1629.
- Campbell, S., Fraser, A., Holliss, B., Schmid, J., & O'Toole, P. W. (1997). Evidence for ethnic tropism of *Helicobacter pylori*. *Infection and Immunity*, 65, 3708–3712.
- Censini, S., Lange, C., Xiang, Z., Crabtree, J.E., Ghiara, P., Borodovsky, M., Rappuoli, R., & Covacci, A. (1996). *cag*, a pathogenicity island of *Helicobacter pylori*, encodes type 1-specific and disease associated virulence factors. *Proceedings of the National Academy of Sciences USA*, 93, 14648–14653.
- Crabtree, J. E., Taylor, J., Heatley, R.V., Shallcross, T. M., Rathbone, B.J., Wyatt, J. I., & Tomkins, D.S. (1991) Mucosal IgA recognition of *Helicobacter pylori* 120 kDa protein, peptic ulceration, and gastric pathology. *Lancet*, 338, 332–335.
- Deguchi, R., Igarashi, M., Watanabe, K., & Takagi, A. (2004). Analysis of the *cag* pathogenicity island and IS605 of *Helicobacter pylori* strains isolated from patients with gastric cancer in Japan. *Alimentary Pharmacology and Therapeutics*, 20, 12–16.
- Fischer, W., Puls, J., Buhrdorf, R., Gebert, B., Odenbreit, S., & Haas, R. (2001). Systematic mutagenesis of the *Helicobacter pylori* *cag* pathogenicity island: essential genes for CagA translocation in host cells and induction of interleukin-8. *Molecular Microbiology*, 42, 1337–1348.
- Glocker, E., Lange, C., Covacci, A., Bereswill, S., Kist, M., & Pahl, H. L. (1998). Proteins encoded by the *cag* pathogenicity island of *Helicobacter pylori* are required for NF- κ B activation. *Infection and Immunity*, 66, 2346–2348.
- Goh, K.L., Cheah, P. L., Noorfaridah, M., Quek, K.F., & Parasakthi, N. (2007). Ethnicity and *Helicobacter pylori* as risk factors for gastric cancer in Malaysia: a prospective case-control study. *American Journal of Gastroenterology*, 102, 40–45.
- Goh, K. L. (1997). Prevalence of and risk factors for *Helicobacter pylori* infection in a multi-racial dyspeptic Malaysian population undergoing endoscopy. *Journal of Gastroenterology and Hepatology*, 12, S29–S35.
- Haron, A., Mazlam, M.Z., Aminuddin, A., Isa, M. R., Madhav, V. K., & Che Ghani, S. (1994). Six year review of gastric carcinoma at the Universiti Kebangsaan Malaysia. *Journal Perubatan UKM*, 16, 13–18.
- Higashi, H., Tsutsumi, R., Fujita, A., Yamazaki, S., Asaka, M., Azuma, T., & Hatakeyama, M. (2002a). Biological activity of the *Helicobacter pylori* virulence factor CagA is determined by variation in the tyrosine phosphorylation sites. *Proceedings of the National Academy of Sciences USA*, 99, 14428–14433.
- Hsu, P. I., Hwang, I. R., Cittelly, D., Lai, K. H., El-Zimaity, H. M., Gutierrez, O., Kim, J. G., Osato, M. S., Graham, D. Y., & Yamaoka, Y. (2002). Clinical presentation in relation to diversity within the *Helicobacter pylori* *cag* pathogenicity island. *American Journal of Gastroenterology*, 97, 2231–2238.

- Hu, P. J., Li, Y. Y., Zhou, M. H., Chen, M. H., Du, G. G., Huang, B. J., Mitchell, H. M., & Hazel, S. L. (1995). *Helicobacter pylori* associated with a high prevalence of duodenal ulcer disease and a low prevalence of gastric cancer in a developing nation. *Gut*, 36, 198–202.
- Ikenoue, T., Maeda, S., Ogura, K., Akanuma, M., Mitsuno, Y., Imai, Y., Yoshida, H., Shiratori, Y., & Omata, M. (2001) Determination of *Helicobacter pylori* virulence by simple gene analysis of the *cag* pathogenicity island. *Clinical and Diagnostic Laboratory Immunology*, 8, 181–186.
- Kang, J. Y., Wee, A., Math, M. V., Guan, R., Tay, H. H., Yap, I., & Sutherland, I. H. (1990). *Helicobacter pylori* and gastritis in patients with peptic ulcer and non-ulcer dyspepsia: ethnic differences in Singapore. *Gut*, 31, 850–853.
- Maeda, S., Yoshida, H., Ikenoue, T., Ogura, K., Kanai, F., Kato, N., Shiratori, Y., & Omata, M. (1999). Structure of *cag* pathogenicity island in Japanese *Helicobacter pylori* isolates. *Gut*, 44, 336–341.
- Maeda, S., Ogura, K., Yoshida, H., Kanai, F., Ikenoue, T., Kato, N., Shiratori, Y., & Omata, M. (1998). Major virulence factors, *VacA* and *CagA*, are commonly positive in *Helicobacter pylori* isolates in Japan. *Gut*, 42, 338–343.
- Mahadeva, S., Raman, M. C., Ford, A. C., Follows, M., Axon, A. T., Goh, K. L., & Moayyedi, P. (2005). Gastroesophageal reflux is more prevalent in western dyspeptics; a prospective comparison of British and South East Asian patients with dyspepsia. *Alimentary Pharmacology and Therapeutics*, 21, 1483–1490.
- Mattar, R., Marques, S. B., Monteiro Mdo, S., Dos Santos, A. F., Iriya, K., & Carrilho, F. J. (2007). *Helicobacter pylori cag* pathogenicity island genes: clinical relevance for peptic ulcer disease development in Brazil. *Journal of Medical Microbiology*, 56, 9–14.
- Mohamed, R., Hanafiah, A., Rose, I. M., Manaf, M. R., Abdullah, S. A., Sagap, I., van Belkum, A., & Yaacob, J. A. (2009). *Helicobacter pylori cagA* gene variants in Malaysians of different ethnicity. *European Journal of Clinical Microbiology and Infectious Diseases*, 28, 865–869.
- Owen, R. J., Peters, T. M., Varea, R., Teare, E., & Saverymuttu, S. (2001). Molecular epidemiology of *Helicobacter pylori* in England: prevalence of *cag* pathogenicity island markers and *IS605* presence in relation to patient age and severity of gastric disease. *FEMS Immunology and Medical Microbiology*, 30, 65–71.
- Peterson, W. L. (1992). *Helicobacter* and peptic ulcer diseases. *The New England Journal of Medicine*, 324, 1043–1048.
- Proença Mòdena, J. L., Lopes Sales, A. I., Olszanski Acrani, G., Russo, R., Vilela Ribeiro, M. A., Fukuhara, Y., da Silveira, W. D., Mòdena, J. L., de Oliveira, R. B., & Brocchi, M. (2007). Association between *Helicobacter pylori* genotypes and gastric disorders in relation to the *cag* pathogenicity island. *Diagnostic Microbiology and Infectious Disease*, 59, 7–16.
- Raj, S. M., Yap, K., Haq, J. A., Singh, S., Hamid, A. (2001). Further evidence for an exceptionally low prevalence of *Helicobacter pylori* infection among peptic ulcer patients in north eastern peninsular Malaysian. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 95, 24–27.
- Ramelah, M., Aminuddin, A., Alfizah, H., Isa, M.R., Jasmi, A. Y., Tan, H. J., Rahman, A. J., Rizal, A. M., & Mazlam, M. Z. (2005). *cagA* gene variants in Malaysian *Helicobacter pylori* strains isolated from patients of different ethnic groups. *FEMS Immunology and Medical Microbiology*, 44, 239–242.
- Segal, E. D., Cha, J., Lo, J., Falkow, S., & Tompkins, L.S. (1999). Altered states: involvement of phosphorylated *CagA* in the induction of host cellular growth changes by *Helicobacter pylori*. *Proceedings of the National Academy of Sciences USA*, 96, 14559–14564.

- Suerbaum, S., & Michetti, P. (2002) *Helicobacter pylori* infection. *The New England Journal of Medicine*, 347, 1175–1186.
- Sheu, S. M., Sheu, B. S., Yang Li, H. B. C., Chu, T. C., & Wu, J. J. (2002). Presence of *iceA1* but not *cagA*, *cagC*, *cagE*, *cagF*, *cagN*, *cagT* or *orf13* genes of *Helicobacter pylori* is associated with more severe gastric inflammation in Taiwanese. *Journal of the Formosan Medical Association*, 10, 18–23.
- Tan, H. J., Rizal, A. M., Rosmadi, M. Y., & Goh, K. L. (2005). Distribution of *Helicobacter pylori* *cagA*, *cagE* and *vacA* in different ethnic groups in Kuala Lumpur, Malaysia. *Journal of Gastroenterology and Hepatology*, 20, 589–594.
- Tan, H. J., & Goh, K. L. (2008). Changing epidemiology of *Helicobacter pylori* in Asia. *Journal of Digestive Diseases*, 9, 186–189.
- van Doorn, L. J., Figueiredo, C., Sanna, R., Plaisier, A., Schneeberger, P., de Boer, W., & Quint, W. (1998). Clinical relevance of the *cagA*, *vacA*, and *iceA* status of *Helicobacter pylori*. *Gastroenterology*, 115, 58–66.
- Xiang, Z., Bugnoli, M., Ponzetto, A., Morgando, A., Figura, N., Covacci, A., Petracca, R., Pennatini, C., Censini, S., & Armellini, D. (1993). Detection in an enzyme immunoassay of an immune response to a recombinant fragment of the 128 kDa protein (*CagA*) of *Helicobacter pylori*. *European Journal of Clinical Microbiology and Infectious Diseases*, 12, 739–745.
- Zheng, P. Y., Hua, J., Yeoh, K. G., & Ho, B. (2000). Association of peptic ulcer with increased expression of Lewis antigens but not *cagA*, *iceA*, and *vacA* in *Helicobacter pylori* isolates in an Asian population. *Gut*, 47, 18–22.