

Review Article

Signaling Behaviour of Abscisic Acid on Physiological Activities in Plants Under Stress

Naimah, N. and Jahan, M. S.*

Department of Agriculture and Biotechnology, Faculty of Bioresources and Food Industry, University Sultan Zainal Abidin, 22200 Besut, Terengganu, Malaysia

ABSTRACT

Plants manage water stress conditions through a variety of signalling networks, which maintain physiological balances of the plants. The phytohormone, abscisic acid (ABA), regulates growth and development of plants by acting on different physiological features of plants. Various stress factors including drought result in raising the endogenous ABA concentration in plant cells. Abscisic acid modifies gene function and its transformation in plants when plants are experiencing stress conditions. Molecular mechanisms show signalling functions of ABA in plants. Several researchers have demonstrated the different roles of ABA in plants such as regulation of stomatal aperture, production of secondary metabolites and signalling cross-link with other molecules. Researchers also show significant interest in and place importance on the signalling regulation of ABA on the growth and physiology of plants. Therefore, this review highlights the signalling pathway of ABA with or without other molecules to restrain growth and physiological parameters of plants under climate change conditions.

Keywords: Arabidopsis, glutathione, guard cells, plant growth, reactive oxygen species, signalling pathway, stomatal aperture, stress factors

ARTICLE INFO

Article history:

Received: 15 May 2017

Accepted: 04 July 2017

E-mail addresses:

noornaimahsyafiqah@gmail.com (Naimah, N.),

sarwarjahan@unisza.edu.my (Jahan, M. S.)

* Corresponding author

INTRODUCTION

Abscisic acid (ABA) is recognised under stress conditions by plants, allowing it to amend physiological features of plants under stress to cope with multiple detrimental environmental indicators that challenge to abiotic and biotic

consequences due to climate change. Currently, biochemical, molecular, genetic and genomic methodologies have shown the fundamental character of the ABA signalling network in plants. Reports on ABA receptors, especially, have further clarified understanding of the functional mechanisms of ABA in plants (Zhang, 2014). Additionally and more importantly, a number of ABA-related genes has been engineered for improvement of stress resistance to provide chances for the progress of new varieties with superior stress resistance. ABA regulates physiological processes, primarily biosynthesis de novo ABA, by relevant enzymes rather than the relocation of remaining pools of ABA (Hartung et al., 2002). Genetic analysis has revealed that different physiological processes such as seed germination, osmotic stress and transpiration are linked to the ABA activity, emphasising the responsibility of ABA in signalling networking in plants (Seo & Koshiba, 2002).

In the last 30 years, several reports confirmed that ABA regulates the stomatal aperture of guard cells. Recent reports have stated that ABA functions with several factors to modulate the growth of plants (Jahan et al., 2008; Khokon et al., 2011; Okuma et al., 2011; Jahan et al., 2016). The status of water maintains ABA strength in the xylem stream. ABA-induced stomatal closure controls physiological functions such as photosynthesis (Davies et al., 1993), which may affect growth, physiology and development of stressed

plants (Jahan et al., 2013; Khairi et al., 2015; Khairi et al., 2016). This review cites new understanding of the current roles of ABA to up-regulate plant physiological functions under climate change conditions for sustainable growth and production of plants.

Discovery and Catabolism of Abscisic Acid

Abscisic acid is a small molecule, which encompasses various functional groups. Addicott et al. (1963) isolated two liable compounds aimed at abscission fruit (cotton) i.e. abscisin I and abscisin II. Abscisic acid was formerly called 'abscisin II'. It plays a significant role in abscission of fruit and bud dormancy and has inhibitory roles as well as promotes physiological functions in plants (Salisbury & Ross, 1992). Stability in biosynthesis and catabolism controls endogenous ABA levels in numerous plants. ABA is a compound that naturally synthesises in plants via a direct pathway, the mevalonic pathway, in chloroplast and via an indirect pathway, the methylerythritol phosphate (MEP) pathway, which appears to occur in cyanobacteria and eukaryotes (Schwartz & Zeevaart, 2010). Stress, for example drought and temperature, highlights the biosynthesis of ABA in plants. In the direct pathway, isomerisation of allofarnesene and oxidation synthesise ABA in fungi (Inomata et al., 2004). In the indirect pathway, carotenoids, chloroplast-produced pigments, which have 40 carbon molecules, appear in the ABA structure.

In the reaction centre, a molecule of violaxanthin was seen in research to produce a molecule of xanthin, which is unstable and therefore, transforms to ABA aldehyde (ABAld). Further oxidation methods used in research resulted in activation of ABA by two enzymes, abscisic aldehyde oxidase and molybdenum cofactor sulfurase. In wilt tomato mutants, *flacca* and *sitiens*, the transformation of ABAld to ABA was constructed in cell-free extracts due to oxidase enzyme (Taylor et al., 1988), which produces ABA (Leydecker et al., 1995). In several plant species, the hydroxylation phase in the ABA catabolic route in plants catalysed C-8' of ABA and produced unstable 8'-hydroxy-ABA (Walton, 1983).

Transportation and Distribution of ABA

Water deficit synthesises ABA in cells sited across the vasculature; it is translocated to the guard cells and stimulates stomatal closure. In this relation, ABA exporter genes and importer genes are expressed; they determine the amount of mobile ABA present in the apoplastic space by balancing export and import of ABA content out of and into vascular tissue (Seo & Koshiba, 2002). ABA is capable of moving both up and down along the stem through the xylem and phloem tissues and parenchyma cells in plants, which do not show polarity with other growth regulators such as auxins (Salisbury & Ross, 1992). Abscisic acid is synthesised in leaves and roots and freely moves from plant to soil

as well as from soil to plant. Roots grow well under soil solution, which contains plants hormones. If the ABA is not at equilibrium, ABA transport from root to soil is disrupted, which also disrupts the signalling relationship between the roots and the shoot (Sauter et al., 2001). Translocation of ABA from cell-to-cell is done through the plasmodesmata and transmembrane transport, after which it is synthesised in the cells. In addition, ABA presences in xylem sap is exported to the outside of the cells (Hartung et al., 2002). After that, ABA is moved to other cells to stimulate physiological reactions in plants.

A plant may uptake ABA from acidic soil, which increases external take-up of ABA concentration by the roots (Freundl et al., 2000). In contrast, roots lose ABA content under high soil pH condition (Degenhardt et al., 2000). This is due to the effects of the reduction of ABA content in the surrounding medium (Freundl et al., 2000). Several factors other than water stress such as salt content, phosphorus deficiency and ammonium increment stimulate ABA biosynthesis in the root but nitrate deficiency and alkaline reduces it (Wolf et al., 1990; Jaschke et al., 1997; Freundl et al., 2000). Therefore, water stress and salt accumulate a higher concentration of ABA in the roots of plants (Sauter et al., 2001). The increment of ABA in roots may increase the flow of ABA in the cells of roots through the symplastic and apoplastic pathway under stress conditions, then translocate to the

guard cells to close the stoma (Sauter et al., 2001; Jahan et al., 2016). In contrast, increasing water flow rapidly decreases ABA concentration through bypass flow of ABA to the endodermis, leading to ABA homeostasis in the xylem (Tardieu et al., 1992; Freundl et al., 1998).

The pH of cells affects the delivery of ABA. ABA biosynthesis occurs in the cytosol when the pH level is at 7.2. Therefore, most of the ABA is ionised in the cytosol and exists as ABA⁻, followed by ABA efflux. Then, the ABA moves through the apoplastic space (Kramer, 2006). At this stage, the pH in the apoplast remains at 5.0-6.0. This pH gradient condition suggests active transport mechanism of apoplastic ABA in cells. However, stomata are closed due to a flux of ABA in well-watered conditions by changing acidic pH to alkaline pH conditions. A little alkalinisation in xylem sap may result in stomatal closure (Wilkinson & Davies, 1997) and might influence leaf growth (Bacon et al., 1998). Apoplastic pH rises to about 7.0 when plants are under water stress conditions, which enables a moderately large percentage of ABA-H to move over extended distances across the apoplast (Wilkinson & Davies, 2002). The transportation of ABA is required to induce ABA signalling under stress conditions. The distribution of ABA is a concern to environmental conditions. Under normal circumstances, ABA concentration in leaves is higher than in roots and stems (Bahrun et al., 2002).

Intercellular Transport of ABA to Stomatal Movement

ABA is a key controller of stomatal movement in guard cells under water scarcity. Several studies have focussed on the signal transduction pathways underlying the ABA-regulated stomatal aperture. ABA is primarily localised in chloroplast and vascular tissue. From here, it is transported to sites of action in guard cells during water shortage (Seo & Koshiba, 2002). Drought-induced pH changes in different parts of the leaves of a plant (Islam et al., 2011) control a complex distribution of ABA throughout the various cells of the leaves. Then ABA gathers in guard cells and persuades stomatal closure (Jahan et al., 2016). Water stress leads to an increment of ABA content in the apoplast but not in the mesophyll cells (Wang & Jia, 1995). Interestingly, ABA translocates from mesophyll and vascular cells to the guard cells to close stomata under stress conditions, but under non-stress conditions, stomatal movement is unaffected by the ABA although both mesophyll and vascular cells contain plentiful ABA content. Drought stress increases ABA content in the mesophyll and vascular cells of leaves and ABA-induced secondary signalling metabolites (Jahan et al., 2016; Munemasa et al., 2007). More prominently, ABA build-up in guard cells seems to protect plants against drought stress (Okuma et al., 2011).

Cross Talk Between ABA and Methyl Jasmonate Acid in Signalling Pathways

Guard cells are specialised cells that control gas exchange and water movement to respond to numerous types of stress due to climate change through the closing and opening of stomatal pores (Kim et al., 2010; Okuma et al., 2011; Jahan et al., 2014). Plants synthesise ABA in response to consequences of climate change and prove the transduction of ABA to guard cells through production of second messengers, for example, cytosolic Ca²⁺, nitric oxide (NO) and reactive oxygen species (ROS) in guard cells (Murata et al., 2015). A plant hormone, jasmonate, regulates stomatal opening in reaction to diverse stimuli such as senescence, wounding and pathogen attacks in plants (Turner et al., 2002). MeJA stimulates ROS intervention in guard cells like ABA under different climate change conditions such as water stress, wounding, salinity and insect attacks (Tsonev et al., 1998; Suhita et al., 2004). Similarities observed in different stomatal aperture experiments involved the NAD(P)H oxidase inhibitor, diphenyleneiodonium (Tsonev et al., 1998). Two NAD(P)H oxidases, AtrbohD and AtrbohF, link methyl jasmonate acid (MeJA) and ROS for signalling in guard cells, proving that MeJA and ABA function in the similar signalling way to persuade stomatal closure. Furthermore, both ABA and MeJA stimulate NO synthase (NOS) guard cells during NO-induced stomatal closure, which indicates a similar path of ABA and MeJA in the signalling force

of stomatal closure (Desikan et al., 2002; Sasaki-Sekimoto et al., 2005). As a result, MeJA- and ABA-induced stomatal closure is due to the production of ROS and NO in guard cells through the stimulation of Ca²⁺ from intracellular stores to the cytosol (Desikan et al., 2002). MeJA does not induce stomatal closure in the *abi2-1* mutant, an ABA-insensitive mutant; this suggests that a protein phosphatase 2C links with MeJA core components in ABA regulation in receptor complexes (Munemasa et al., 2007; Desikan et al., 2002; Murata et al., 2001). ABA and MeJA signalling might affect ABA receptor developments to control downstream signal mechanisms.

Abscisic Acid Signalling in Glutathione Functions on Guard Cells

Abscisic acid stimulates different physiological processes of a plant in stress conditions (Finkelstein et al., 2002). ABA features on GSH variation in guard cells of Arabidopsis (Okuma et al., 2011; Jahan et al., 2014). Water stress promotes the primary gathering of ABA that generates ROS creation, which might adjust GSH biosynthesis in plants (Murata et al., 2015). ABA signalling in GSH was studied earlier in guard cells of the Arabidopsis plants (Jahan et al., 2008; Jahan et al., 2011; Okuma et al., 2011; Munemasa et al., 2013). The Arabidopsis mutant, GLUTATHIONE S-TRANSFERASE U17 (*AtGSTU17*), stores a greater amount of ABA in the presence of GSH (Cheng et al., 2013). Therefore, the functions of GSH

with ABA in plants would be a critical aspect for the plants' ability to cope with climate change conditions (Cheng et al., 2013). ABA affects GSH content in ratio to GSSG, GR activity and γ -ECS transcript levels in maize genotypes (Okuma et al., 2011; Cheng et al., 2013). Glutathione peroxidase3 (*atgpx3*) controls H₂O₂ homeostasis in ABA-mediated stomatal movement (Miao et al., 2006), which suggests that ABA signalling with GSH is linked to the stomatal movement in guard cells of *Arabidopsis* (Okuma et al. 2011; Jahan et al., 2014). Therefore, GSH-deficient mutants, the *chl-1* and cadmium sensitive mutant (*cad2-1*), exhibit a higher sensitivity to ABA than the wild types (Okuma et al., 2011; Jahan et al., 2011, 2008). In addition, Munemasa et al. (2013) confirm that GSH modulates the signalling compartment of ABA in guard cells.

It has been genetically and chemically confirmed that GSH does not affect ROS production in guard cells as well as the activation of Ca²⁺-permeable channel (ICa) (Akter et al., 2013; Okuma et al., 2011), which suggests that GSH might play a role in coupling H₂O₂ to downstream an ABA-induced signalling cascade in guard cells (Okuma et al. 2011). In a current development, Munemasa et al. (2013) stated that GSH depletion affects ROS signalling in apoplastic space rather than in the cytosol of guard cells during ABA-induced stomatal closure (Munemasa et al., 2013). Different studies have stated a similar result, that apoplastic ROS signals regulate stomatal movement (Hossain

et al., 2013; An et al., 2008). Therefore, GSH-modulated ABA signalling of a GSH-deficient mutant (Jahan et al., 2008) is not conditional to the production of H₂O₂ in guard cells (Munemasa et al., 2013).

Abscisic Acid Signalling on Growth of Plants

Climate change is becoming a severe problem for plant growth in several regions of the world. Plants respond to drought conditions through intensely complicated mechanisms such as genetic and physiological processes (Chaves et al., 2003; Izanloo et al., 2008). Water stress influences plant growth and biomass production remarkably (Chaves et al., 2003; Hisao, 1973). The leaf area of *Leymus chinensis*, for instance, is significantly stimulated by rewatering compared to a short-term drought (Xu & Zhou, 2007). In another instance, tiller production of *Leymus chinensis* recovered after rewatering, suggesting that development of meristems of the tiller might play an important role in response to water stress conditions (Xu & Zhou, 2009). Water stress is closely related to the elevation of ABA in guard cells to close stomata (Jahan et al., 2016), which might reduce gas exchange through the opening of guard cells. Reactive oxygen species (ROS) increases during drought conditions because of the influence of ABA production (Jahan et al., 2008). Water application reduced ROS production in guard cells and the retrieval net photosynthetic and transpiration rate in plants (Okuma et al.,

2011; Khairi et al., 2016). Low relative water content increased ROS production (Murata et al., 2015), which is also related to the ABA signalling for plant growth. Stoma closure due to drought conditions limits photosynthesis in plants (Chaves et al., 2003). Drought reduces light-dependent photosynthesis and stomatal conductance in plants (Peñuelas et al., 2004; Xu & Zhou, 2009). This was consistent with the results obtained by Nozulai et al. (2015) and Khaiti et al. (2016) in rice plants, where reduction of water content reduced plant physiological parameters. This result also suggests that ABA might have a significant role in gas exchange in drought conditions. In addition, Izanloo et al. (2008) stated that water stress declined quantum yield in the photosystem II, indicating a functional activity of ABA in regulating light reaction in plants. A defect of the light-harvesting mutant, *chl-1*, led to higher sensitivity to ABA than was seen in wild types (Jahan et al., 2014, 2016). Therefore, *chl-1* mutant plants are deficient in morphological, physiologic and yield parameters compared to wild types of the plant (Jahan et al., 2014, 2016). It is suggested that ABA controls growth and development of plants through multi-signalling waves.

CONCLUSION

Considering the latest documentation, it is clear that ABA has a distinct role in the hormonal signalling pathway in controlling physiological functions of plants. There is evidence that ABA modulates physiological changes in plants in the presence of stress

conditions through a signalling network with other plant hormones as well as antioxidants. However, understanding the workings of the protocols of the gene that controls ABA function in plants needs further study. Further study and experimental proof are also necessary for understanding the role of ABA in different hormonal signalling cascades for a clearer picture of the ABA signalling-web on the growth and physiology of plants.

ACKNOWLEDGEMENT

This work was funded by a grant, FRGS/2/2014/STWN03/UNISZA/02/1, from the KPT, Kuala Lumpur, Malaysia awarded to M. S. Jahan.

REFERENCES

- Addicott, F. T., Lyon, J. L., Ohkuma, K., Thiessen, W. E., Carns, H. R., Smith, O. E., ... & Wareing, P. F. (1968). Abscisic acid: A new name for abscisin II (dormin). *Science*, *159*(3822), 1493-1493.
- Akter, N., Okuma, E., Sobahan, M. A., Uraji, M., Munemasa, S., Nakamura, Y., ... & Murata, Y. (2013). Negative regulation of methyl jasmonate-induced stomatal closure by glutathione in *Arabidopsis*. *Journal of plant growth regulation*, *32*(1), 208-215.
- An, Z., Jing, W., Liu, Y. L., & Zhang, W. H. (2008). Hydrogen peroxide generated by copper amine oxidase is involved in abscisic acid-induced stomatal closure in *Vicia faba*. *Journal of Experimental Botany*, *59*(4), 815-825.
- Bacon, M. A., Wilkinson, S., & Davies, W. J. (1998). pH-regulated leaf cell expansion in droughted plants is abscisic acid dependent. *Plant physiology*, *118*(4), 1507-1515.

- Bahrn, A., Jensen, C. R., Asch, F., & Mogensen, V. O. (2002). Drought-induced changes in xylem pH, ionic composition, and ABA concentration act as early signals in field-grown maize (*Zea mays L.*). *Journal of Experimental Botany*, 53(367), 251–63.
- Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant response to drought—from genes to the whole plant. *Functional Plant Biology*, 30(3), 239–64.
- Cheng, M. C., Liao, P. M., Kuo, W. W., & Lin T. P. (2013). The *Arabidopsis* ethylene response factor1 regulates abiotic stress-responsive gene expression by binding to different cis-acting elements in response to different stress signals. *Plant Physiology*, 162(3), 1566–1582.
- Davies, W. J., Trejo, C. L., & Palmer, S. J. (1993) ABA and the control of growth and physiology of stressed plants. In M. B. Jackson & C. R. Black (Eds.), *Interacting stresses on plants in a changing climate. NATO ASI series (Series I: Global environmental change)* (vol. 16). Berlin, Heidelberg: Springer.
- Degenhardt, B., Gimmler, H., Hose, E., & Hartung, W. (2000). Effect of alkaline and saline substrates on ABA contents, distribution and transport in plant roots. *Plant and Soil*, 225(1), 83–94.
- Desikan, R., Griffiths, R., Hancock, J., & Neill, S. (2002). A new role of an old enzyme: Nitrate reductase-mediated nitric oxide generation is required for abscisic acid-induced stomatal closure in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences USA*, 99(25), 16314–16318.
- Finkelstein, R. R., Gampala, S. S., & Rock, C. D. (2002). Abscisic acid signaling in seeds and seedlings. *The Plant Cell Online*, 14(suppl 1), S15-S45.
- Freundl, E., Steudle, E., & Hartung, W. (1998). Water uptake by roots of maize and sunflower affects the radial transport of abscisic acid and the ABA concentration in the xylem. *Planta*, 207(1), 8–19.
- Freundl, E., Steudle, E., & Hartung, W. (2000). Apoplastic transport of abscisic acid through roots of maize: Effect of the exodermis. *Planta*, 210(2), 222–231.
- Hartung, W., Sauter, A., & Hose, E. (2002). Abscisic acid in the xylem: where does it come from, where does it go to? *Journal of Experimental Botany*, 53(366), 27–32.
- Hisao, T. C. (1973). Plant responses to water stress. *Annual Review of Plant Physiology*, 24(1), 519–70.
- Hossain, M. S., Ye, W., Hossain, M. A., Okuma, E., Uraji, M., Nakamura, Y., ... & Murata, Y. (2013). Glucosinolate degradation products, isothiocyanates, nitriles, and thiocyanates, induce stomatal closure accompanied by peroxidase-mediated reactive oxygen species production in *Arabidopsis thaliana*. *Bioscience, biotechnology, and biochemistry*, 77(5), 977–983.
- Inomata, M., Hirai, N., Yoshida, R., & Ohigashi, H. (2004). Biosynthesis of abscisic acid by the direct pathway via ionylideneethane in a fungus, *Cercospora cruenta*. *Bioscience Biotechnology Biochemistry*, 68(12), 2571–2580.
- Islam, M. M., Hoque, M. A., Okuma, E., Jannat, R., Banu, M. N. A., Jahan, M. S., ... & Murata, Y. (2009). Proline and glycinebetaine confer cadmium tolerance on tobacco bright yellow-2 cells by increasing ascorbate-glutathione cycle enzyme activities. *Bioscience, biotechnology, and biochemistry*, 73(10), 2320-2323.

- Izanloo, A., Condon, A. G., Langridge, P., Tester, M., & Schnurbusch, T. (2008) Different mechanisms of adaptation to cyclic water stress in two South Australian bread wheat cultivars. *Journal of Experimental Botany*, 59(12), 3327–3346.
- Jahan, M. S., Nakamura, Y., & Murata, Y. (2011). Histochemical quantification of GSH contents in guard cells of *Arabidopsis thaliana*. *Science Asia*, 37, 291–295.
- Jahan, M. S., Nozulaidi, M., Khairi, M., & Mat, N. (2016). Light-harvesting complexes in photosystem II regulate glutathione-induced sensitivity of Arabidopsis guard cells to abscisic acid. *Journal of Plant Physiology*, 195, 1–8.
- Jahan, M. S., Nozulaidi, M., Khandaker, M. M., Afifah, A., & Husna, N. (2014). Control of plant growth and water loss by a lack of light-harvesting complexes in photosystem-II in Arabidopsis thaliana ch1-1 mutant. *Acta Physiologia Plantarum*, 36(7), 1627–1635.
- Jahan, M. S., Nozulaidi, M. B. N., Khairi, M. B. C. L., & Khanif, Y. M. (2013). Effects of water stress on rice production: Bioavailability of potassium in soil. *Journal of Stress Physiology and Biochemistry*, 9(2), 97–107.
- Jahan, M. S., Ogawa, K., Nakamura, Y., Shimoishi, Y., Mori, I. C., & Murata, Y. (2008). Deficient glutathione in guard cells facilitates abscisic acid-induced stomatal closure but does not affect light-induced stomatal opening. *Bioscience Biotechnology Biochemistry*, 72(10), 2795–2798.
- Jaschke, W. D., Peuke, A. D., Pate, J. S., & Hartung, W. (1997). Transport, synthesis and catabolism of abscisic acid (ABA) in intact plants of castor bean (*Ricinus communis* L.) under phosphate deficiency and moderate salinity. *Journal of Experimental Botany*, 48(9), 1737–1747.
- Khairi, M., Nozulaidi, M., Afifah, A., & Jahan, M. S. (2015). Effect of various water regimes on rice production in lowland irrigation. *Australian Journal of Crop Science*, 9(2), 153–159.
- Khairi, M., Nozulaidi, M., & Jahan, M. S. (2016). Effects of flooding and alternate wetting and drying on the yield performance of upland rice. *Pertanika Journal of Tropical Agricultural Science*, 39(3), 299–309.
- Khokon, M. A. R., Jahan, M. S., Rahman, T., Hossain, M. A., Munemasa, S., Mori, I. C., ... & Murata, Y. (2011). Allylisothiocyanate (AITC) induces stomatal closure in Arabidopsis. *Plant Cell Environment*, 34(11), 900–1906.
- Kim, T. H., Böhmer, M., Hu, H., Nishimura, N., & Schroeder, J. I. (2010). Guard cell signal transduction network: Advances in understanding abscisic acid, CO₂ and Ca²⁺ signaling. *Annual Review of Plant Biology*, 61, 561–591.
- Kramer, E. M. (2006). How far can a molecule of weak acid travel in the apoplast or xylem? *Plant physiology*, 141(4), 1233–1236.
- Leydecker, M. T., Moureaux, T., Kraepiel, Y., Schnorr, K., & Caboche, M. (1995). Molybdenum cofactor mutants, specifically impaired in xanthine dehydrogenase activity and abscisic acid biosynthesis, simultaneously overexpress nitrate reductase. *Plant physiology*, 107(4), 1427–1431.
- Miao, Y., Lv, D., Wang, P., Wang, X. C., Chen, J., Miao, C., & Song, C. P. (2006). An Arabidopsis glutathione peroxidase functions as both a redox transducer and a scavenger in abscisic acid and drought stress responses. *Plant Cell*, 18(10), 2749–2766.

- Munemasa, S., Muroyama, D., Nagahashi, H., Nakamura, Y., Mori, I. C., & Murata, Y. (2013). Regulation of reactive oxygen species-mediated abscisic acid signaling in guard cells and drought tolerance by glutathione. *Frontier Plant Science*, 4, 472–477.
- Munemasa, S., Oda, K., Watanabe-Sugimoto, M., Nakamura, Y., Shimoiishi, Y., & Murata, Y. (2007). The *coronatine-insensitive 1* mutation reveals the hormonal signaling interaction between abscisic acid and methyl jasmonate in *Arabidopsis* guard cells. Specific impairment of ion channel activation and second messenger production. *Plant physiology*, 143(3), 1398–1407.
- Murata, Y., Mori, I. C., & Munemasa, S. (2015). Diverse stomatal signaling and the signal integration mechanism. *Annual Review of Plant Biology*, 66, 369–392.
- Murata, Y., Pei, Z. M., Mori, I. C., & Schroeder, J. (2001). Abscisic acid activation of plasma membrane Ca²⁺ channels in guard cells requires cytosolic NAD(P)H and is differentially disrupted upstream and downstream of reactive oxygen species production in *abi1-1* and *abi2-1* protein phosphatase2C mutants. *Plant Cell*, 13(11), 2513–2523.
- Nozulaidi, M., Khairi, M., & Jahan, M. S. (2015). Effects of different salinity levels on rice production. *Australian Journal of Basic and Applied Sciences*, 9, 524–530.
- Okuma, E., Jahan, M. S., Munemasa, S., Ogawa, K., Watanabe-Sugimoto, M., Nakamura, Y., ... & Murata, Y. (2011). Negative regulation of abscisic acid-induced stomatal closure by glutathione in *Arabidopsis*. *Journal of Plant Physiology*, 168(17), 2048–2055.
- Peñuelas, J., Munné-Bosch, S., Llusià, J., & Filella, I. (2004). Leaf reflectance and photo- and antioxidant protection in field-grown summer-stressed *Phillyrea angustifolia*. Optical signals of oxidative stress? *New phytology*, 162(1), 115–124.
- Salisbury, F. B., & Ross, C.W. (1992). *Plant physiology*. Belmont, CA: Wadsworth.
- Sasaki-Sekimoto, Y., Taki, N., Obayashi, T., Aono, M., et al. (2005). Coordinated activation of metabolic pathways for antioxidants and defense compounds by jasmonates and their roles in stress tolerance in *Arabidopsis*. *Plant Journal*, 44(4), 653–668.
- Sauter, A., Davies, W. J., & Hartung, W. (2001). The long-distance abscisic acid signal in the droughted plants: The fate of the hormone on its way from root to shoot. *Journal of Experimental Botany*, 52(363), 1991–1997.
- Schwartz, S. H., & Zeevaart, J. A. D. (2010). Abscisic acid biosynthesis and metabolism. In P. J. Davies (Ed.), *Plant hormones* (3rd Ed.) (pp. 137–55). New York: Springer.
- Seo, M., & Koshiba, T. (2002). Complex regulation of ABA biosynthesis in plants. *Trends Plant Science*, 7(1), 41–48.
- Suhita, D., Raghavendra, A. S., Kwak, J. M., & Vavasseur, A. (2004) Cytoplasmic alkalization precedes reactive oxygen species production during methyl jasmonate- and abscisic acid-induced stomatal closure. *Plant Physiology*, 134(4), 1536–1545.
- Tardieu, F., Zhang, J., Katerji, N., Bethenod, O., Palmer, S., & Davies, W. J. (1992). Xylem ABA controls the stomatal conductance of field-grown maize subjected to soil compaction or drying soil. *Plant Cell and Environment*, 15(2), 193–197.
- Taylor, I. B., Linforth, R. S. T., Alnaieb, R. J., Bowman, W. R., & Marples, B. A. (1988). The wilted tomato mutants Flacca and Sitiens are impaired in the oxidation of Aba-Aldehyde to ABA. *Plant and Cell Environment*, 11(8), 739–45.
- Tsonev, T. D., Lazova, G. N., Stoinova, Z. G., & Popova, L. P. (1998). A possible role for jasmonic acid in adaptation of barley seedlings to salinity stress. *Journal of Plant Growth Regulation*, 17(3), 153–159.

- Turner, J. G., Ellis, C., & Devoto, A. (2002). The jasmonate signal pathway. *Plant Cell*, 14(Suppl 1), 153–164.
- Walton, D. C. (1983). Structure-activity relationships of abscisic acid analogs and metabolites. In F. T. Addicott (Ed.), *Abscisic acid* (p. 113–46). New York: Praeger.
- Wang, X. C., & Jia, W. (1995). Stress-induced stomatal closure in relation to compartmentation and redistribution of ABA in intact leaves of *Vicia faba* L. *Acta Phytophysiologica Sinica*, 21(4), 324–328.
- Wilkinson, S., & Davies, W. J. (1997). Xylem sap pH increase: A drought signal received at the apoplastic face of the guard cell that involves the suppression of saturable abscisic acid uptake by the epidermal symplast. *Plant Physiology*, 113(2), 559–573.
- Wilkinson, S., & Davies, W. J. (2002). ABA-based chemical signalling: The co-ordination of responses to stress in plants. *Plant and Cell Environment*, 25(2), 195–210.
- Wolf, O., Jeschke, W. D., & Hartung, W. (1990). Long-distance transport of abscisic acid in NaCl-treated intact plants of *Lupinus albus*. *Journal of Experimental Botany*, 41(5), 593–600.
- Xu, Z. Z., & Zhou, G. S. (2007). Photosynthetic recovery of a perennial grass *Leymus chinensis* after different periods of soil drought. *Plant Production Science*, 10(3), 277–285.
- Xu, Z. Z., & Zhou, G. S. (2009). Shimizu H. Are plant growth and photosynthesis limited by pre-drought following rewatering in grass? *Journal of Experimental Botany*, 60(13), 3737–49.
- Zhang, D. P. (Ed.). (2014). *Abscisic acid: Metabolism, transport and signaling*. New York, NY: Springer Dordrecht Heidelberg.

